

INNER AND OUTER AUTOMORPHISMS OF FREE METABELIAN NILPOTENT LIE ALGEBRAS

VESSELIN DRENSKY AND ŞEHMUS FINDIK

ABSTRACT. Let $L_{m,c}$ be the free metabelian nilpotent of class c Lie algebra of rank m over a field K of characteristic 0. We describe the groups of inner and outer automorphisms of $L_{m,c}$. To obtain this result we first describe the groups of inner and continuous outer automorphisms of the completion $\widehat{F}_m$ with respect to the formal power series topology of the free metabelian Lie algebra F_m of rank m .

INTRODUCTION

Let L_m be the free Lie algebra of rank $m \geq 2$ over a field K of characteristic 0 with free generators $x_1, \dots, x_m$ and let $L_{m,c} = L_m / (L_m'' + L_m^{c+1})$ be the free metabelian nilpotent of class c Lie algebra. This is the relatively free algebra of rank m in the variety of Lie algebras $\mathfrak{A}^2 \cap \mathfrak{N}_c$, where $\mathfrak{A}^2$ is the metabelian (solvable of class 2) variety of Lie algebras and $\mathfrak{N}_c$ is the variety of all nilpotent Lie algebras of class at most c .

The initial goal of our paper was to describe the groups of inner automorphisms $\text{Inn}(L_{m,c})$ and outer automorphisms $\text{Out}(L_{m,c})$ of the Lie algebra $L_{m,c}$. The automorphism group $\text{Aut}(L_{m,c})$ is a semidirect product of the normal subgroup $\text{IA}(L_{m,c})$ of the automorphisms which induce the identity map modulo the commutator ideal of $L_{m,c}$ and the general linear group $\text{GL}_m(K)$. Since $\text{Inn}(L_{m,c}) \subset \text{IA}(L_{m,c})$, for the description of the factor group $\text{Out}(L_{m,c}) = \text{Aut}(L_{m,c}) / \text{Inn}(L_{m,c})$ it is sufficient to know only $\text{IA}(L_{m,c}) / \text{Inn}(L_{m,c})$.

The composition of two inner automorphisms is an inner automorphism obtained by the Baker-Campbell-Hausdorff formula which gives the solution z of the equation $e^z = e^x \cdot e^y$ for non-commuting x and y . If x, y are the generators of the free associative algebra $A = K\langle x, y \rangle$, then z is a formal power series in the completion $\widehat{A}$ with respect to the formal power series topology. The homogeneous components of z are Lie elements. Hence, it is natural to work in the completion $\widehat{F}_m$ with respect to the formal power series topology of the free metabelian Lie algebra $F_m = L_m / L_m''$ and to study the groups of its inner automorphisms and of its continuous outer automorphisms. Then the results for $\text{Inn}(L_{m,c})$ and $\text{Out}(L_{m,c})$ follow easily by factorization modulo $(\widehat{F}_m)^{c+1}$. Gerritzen [G] found a simple version of the Baker-Campbell-Hausdorff formula when applied to the algebra $\widehat{F}_2$. This provides a nice expression of the composition of inner automorphisms in $\widehat{F}_m$.

2010 *Mathematics Subject Classification.* 17B01, 17B30, 17B40.

Key words and phrases. free metabelian Lie algebras, inner automorphisms, outer automorphisms, Baker-Campbell-Hausdorff formula.

The research of the second named author was partially supported by the Council of Higher Education (YÖK) in Turkey.

A result of Shmel'kin [Sh] states that the free metabelian Lie algebra F_m can be embedded into the abelian wreath product $A_m \text{wr} B_m$, where A_m and B_m are m -dimensional abelian Lie algebras with bases $\{a_1, \dots, a_m\}$ and $\{b_1, \dots, b_m\}$, respectively. The elements of $A_m \text{wr} B_m$ are of the form $\sum_{i=1}^m a_i f_i(t_1, \dots, t_m) + \sum_{i=1}^m \beta_i b_i$, where f_i are polynomials in $K[t_1, \dots, t_m]$ and $\beta_i \in K$. This allows to introduce partial derivatives in F_m with values in $K[t_1, \dots, t_m]$ and the Jacobian matrix $J(\phi)$ of an endomorphism ϕ of F_m . Restricted on the semigroup $\text{IE}(F_m)$ of endomorphisms of F_m which are identical modulo the commutator ideal F'_m , the map $J : \phi \rightarrow J(\phi)$ is a semigroup monomorphism of $\text{IE}(F_m)$ into the multiplicative semigroup of the algebra $M_m(K[t_1, \dots, t_m])$ of $m \times m$ matrices with entries from $K[t_1, \dots, t_m]$. We give the explicit form of the Jacobian matrices of inner automorphisms of $\widehat{F_m}$ and of the coset representatives of the continuous outer automorphisms in $\text{IOut}(\widehat{F_m})$.

Finally we transfer the obtained results to the algebra $L_{m,c}$ and obtain the description of $\text{Inn}(L_{m,c})$ and $\text{IOut}(L_{m,c})$.

1. PRELIMINARIES

Let L_m be the free Lie algebra of rank $m \geq 2$ over a field K of characteristic 0 with free generators $x_1, \dots, x_m$ and let $L_{m,c} = L_m / (L''_m + L_m^{c+1})$ be the free metabelian nilpotent of class c Lie algebra. It is freely generated by $y_1, \dots, y_m$, where $y_i = x_i + (L''_m + L_m^{c+1})$, $i = 1, \dots, m$. We use the commutator notation for the Lie multiplication. Our commutators are left normed:

$$[u_1, \dots, u_{n-1}, u_n] = [[u_1, \dots, u_{n-1}], u_n], \quad n = 3, 4, \dots$$

In particular,

$$L_m^k = \underbrace{[L_m, \dots, L_m]}_{k \text{ times}}.$$

For each $v \in L_{m,c}$, the linear operator $\text{adv} : L_{m,c} \rightarrow L_{m,c}$ defined by

$$u(\text{adv}) = [u, v], \quad u \in L_{m,c},$$

is a derivation of $L_{m,c}$ which is nilpotent and $\text{ad}^c v = 0$ because $L_{m,c}^{c+1} = 0$. Hence the linear operator

$$\exp(\text{adv}) = 1 + \frac{\text{adv}}{1!} + \frac{\text{ad}^2 v}{2!} + \dots = 1 + \frac{\text{adv}}{1!} + \frac{\text{ad}^2 v}{2!} + \dots + \frac{\text{ad}^{c-1} v}{(c-1)!}$$

is well defined and is an automorphism of $L_{m,c}$. The set of all such automorphisms forms a normal subgroup of the group of all automorphisms $\text{Aut}(L_{m,c})$ of $L_{m,c}$. This group is called the inner automorphism group of $L_{m,c}$ and is denoted by $\text{Inn}(L_{m,c})$. The factor group $\text{Aut}(L_{m,c})/\text{Inn}(L_{m,c})$ is called the outer automorphism group of $L_{m,c}$ and is denoted by $\text{Out}(L_{m,c})$. The automorphism group $\text{Aut}(L_{m,c})$ is a semidirect product of the normal subgroup $\text{IA}(L_{m,c})$ of the automorphisms which induce the identity map modulo the commutator ideal of $L_{m,c}$ and the general linear group $\text{GL}_m(K)$. Since $\text{Inn}(L_{m,c}) \subset \text{IA}(L_{m,c})$, for the description of $\text{Out}(L_{m,c}) = \text{Aut}(L_{m,c})/\text{Inn}(L_{m,c})$ it is sufficient to know only $\text{IA}(L_{m,c})/\text{Inn}(L_{m,c})$.

Let R be an algebra over a field K of characteristic 0. We consider the topology on R induced by the series $R \supseteq R^2 \supseteq R^3 \supseteq \dots$. This is the topology in which the sets $r + R^k$, $r \in R$, $k \geq 1$, form a basis for the open sets. It is called the *formal power series topology* on R .

Let $F_m = L_m/L_m''$ be the free metabelian Lie algebra of rank m . We shall denote the free generators of F_m with the same symbols $y_1, \dots, y_m$ as the free generators of $L_{m,c}$, but now $y_i = x_i + L_m''$, $i = 1, \dots, m$. It is well known, see e.g. [Ba], that

$$[y_{i_1}, y_{i_2}, y_{i_{\sigma(i_3)}}, \dots, y_{i_{\sigma(k)}}] = [y_{i_1}, y_{i_2}, y_{i_3}, \dots, y_{i_k}],$$

where σ is an arbitrary permutation of $3, \dots, k$ and that F_m' has a basis consisting of all

$$[y_{i_1}, y_{i_2}, y_{i_3}, \dots, y_{i_k}], \quad 1 \leq i_j \leq m, \quad i_1 > i_2 \leq i_3 \leq \dots \leq i_k.$$

Let $(F_m)_{(k)}$ be the subspace of F_m spanned by all monomials of total degree k in $y_1, \dots, y_m$. We consider the formal power series topology on F_m . The completion $\widehat{F_m}$ of F_m with respect to this topology may be identified with the complete direct sum $\bigoplus_{i \geq 1} (F_m)_{(i)}$. The elements $f \in \widehat{F_m}$ are formal power series

$$f = f_1 + f_2 + \dots, \quad f_i \in (F_m)_{(i)}, \quad i = 1, 2, \dots$$

The composition of two inner automorphisms in $\text{Inn}(L_{m,c})$ is also an inner automorphism. It can be obtained by the Baker-Campbell-Hausdorff formula which gives the solution z of the equation $e^z = e^x \cdot e^y$ for non-commuting x and y , see e.g. [Bo] and [Se]. If x, y are the generators of the free associative algebra $A = K\langle x, y \rangle$, then

$$z = x + y + \frac{[x, y]}{2} - \frac{[x, y, x]}{12} + \frac{[x, y, y]}{12} - \frac{[x, y, x, y]}{24} + \dots$$

is a formal power series in the completion $\widehat{A}$ with respect to the formal power series topology. The homogeneous components of z are Lie elements. Hence, studying the inner and the outer automorphisms of $L_{m,c}$, it is convenient to work in the completion $\widehat{F_m}$ and to study the groups of its inner automorphisms and of its continuous outer automorphisms. Working in the algebra $F_2 = L_2/L_2''$ with generators $y_1 = x$, $y_2 = y$, the element $z \in \widehat{F_2}$ in the Baker-Campbell-Hausdorff formula has the form

$$\begin{aligned} z &= x + y + \sum_{a,b \geq 0} c_{ab} [x, y, \underbrace{x, \dots, x}_a, \underbrace{y, \dots, y}_b] = x + y + \sum_{a,b \geq 0} c_{ab} [x, y] \text{ad}^a x \text{ad}^b y \\ &= x + y + [x, y] c(\text{ad} x, \text{ad} y), \end{aligned}$$

where

$$c(t, u) = \sum_{a,b \geq 0} c_{ab} t^a u^b \in \mathbb{Q}[[t, u]].$$

Gerritzen [G] found a nice description of the formal power series in commuting variables $c(t, u)$ which corresponds to $z \in \widehat{F_2}$. Further we shall use it to obtain an expression of the composition of inner automorphisms in $\widehat{F_m}$. We present a slightly modified version of the result of Gerritzen due to the fact that he uses right normed commutators (and not left normed commutators as we do). Recall that F_2 is isomorphic to the Lie subalgebra generated by x and y in the associative algebra A/C^2 , where $C = A[A, A]A$ is the commutator ideal of $A = K\langle x, y \rangle$.

Proposition 1. (Gerritzen [G]) *If $z \in \widehat{F_2}$ is the solution of the equation $e^z = e^x \cdot e^y$ in the algebra $\widehat{A/C^2}$, then*

$$z = x + y + [x, y] c(\text{ad} x, \text{ad} y),$$

where

$$c(t, u) = \frac{e^u h(t) - h(u)}{e^{t+u} - 1} \in K[[t, u]], \quad h(v) = \frac{e^v - 1}{v} \in K[[v]].$$

Proof. Following Gerritzen [G], if D_v is the derivation of the free metabelian algebra F_2 defined by $D_v(u) = [v, u]$, $u \in F_2$, then the solution $z \in \widehat{F_2}$ of the equation $e^z = e^x \cdot e^y$ has the form

$$z = x + y + H_0(D_x, D_y)[x, y],$$

where

$$H_0(t, u) = \frac{h(-t) - h(u)}{e^{-t} - e^u}, \quad h(v) = \frac{e^v - 1}{v}.$$

Since $D_x = -\text{adx}$ and $D_y = -\text{ady}$, we obtain

$$z = x + y + H_0(D_x, D_y)[x, y] = x + y + [x, y]H_0(-\text{adx}, -\text{ady}) = x + y + [x, y]c(\text{adx}, \text{ady}),$$

$$c(t, u) = H_0(-t, -u) = \frac{h(t) - h(-u)}{e^t - e^{-u}} = \frac{e^u h(t) - h(u)}{e^{t+u} - 1}$$

after easy computations. $\square$

Pay attention that both the nominator and the denominator of $c(t, u)$ are divisible by $t + u$. After the cancellation of the common factor $t + u$ the denominator becomes an invertible element of $K[[t, u]]$.

Now we collect the necessary information about wreath products, Jacobian matrices and formal power series. For details and references see e.g. [BD]. Let $K[t_1, \dots, t_m]$ be the (commutative) polynomial algebra over K freely generated by the variables $t_1, \dots, t_m$ and let A_m and B_m be the abelian Lie algebras with bases $\{a_1, \dots, a_m\}$ and $\{b_1, \dots, b_m\}$, respectively. Let C_m be the free right $K[t_1, \dots, t_m]$ -module with free generators $a_1, \dots, a_m$. We give it the structure of a Lie algebra with trivial multiplication. The abelian wreath product $A_m \text{wr} B_m$ is equal to the semidirect sum $C_m \rtimes B_m$. The elements of $A_m \text{wr} B_m$ are of the form $\sum_{i=1}^m a_i f_i(t_1, \dots, t_m) + \sum_{i=1}^m \beta_i b_i$, where f_i are polynomials in $K[t_1, \dots, t_m]$ and $\beta_i \in K$. The multiplication in $A_m \text{wr} B_m$ is defined by

$$[C_m, C_m] = [B_m, B_m] = 0,$$

$$[a_i f_i(t_1, \dots, t_m), b_j] = a_i f_i(t_1, \dots, t_m) t_j, \quad i, j = 1, \dots, m.$$

Hence $A_m \text{wr} B_m$ is a metabelian Lie algebra and every mapping $\{y_1, \dots, y_m\} \rightarrow A_m \text{wr} B_m$ can be extended to a homomorphism $F_m \rightarrow A_m \text{wr} B_m$. As a special case of the embedding theorem of Shmel'kin [Sh], the homomorphism $\varepsilon : F_m \rightarrow A_m \text{wr} B_m$ defined by $\varepsilon(y_i) = a_i + b_i$, $i = 1, \dots, m$, is a monomorphism. If

$$f = \sum [y_i, y_j] f_{ij}(\text{ady}_1, \dots, \text{ady}_m), \quad f_{ij}(t_1, \dots, t_m) \in K[t_1, \dots, t_m],$$

then

$$\varepsilon(f) = \sum (a_i t_j - a_j t_i) f_{ij}(t_1, \dots, t_m).$$

The next lemma follows from [Sh], see also [BD].

Lemma 2. *The element $\sum_{i=1}^m a_i f_i(t_1, \dots, t_m)$ of C_m belongs to $\varepsilon(F'_m)$ if and only if $\sum_{i=1}^m t_i f_i(t_1, \dots, t_m) = 0$.*

The embedding of F_m into $A_m \text{wr} B_m$ allows to introduce partial derivatives in F_m with values in $K[t_1, \dots, t_m]$. If $f \in F_m$ and

$$\varepsilon(f) = \sum_{i=1}^m \beta_i b_i + \sum_{i=1}^m a_i f_i(t_1, \dots, t_m), \quad \beta_i \in K, f_i \in K[t_1, \dots, t_m],$$

then

$$\frac{\partial f}{\partial y_i} = f_i(t_1, \dots, t_m).$$

The Jacobian matrix $J(\phi)$ of an endomorphism ϕ of F_m is defined as

$$J(\phi) = \left(\frac{\partial \phi(y_j)}{\partial y_i} \right) = \begin{pmatrix} \frac{\partial \phi(y_1)}{\partial y_1} & \dots & \frac{\partial \phi(y_m)}{\partial y_1} \\ \vdots & \ddots & \vdots \\ \frac{\partial \phi(y_1)}{\partial y_m} & \dots & \frac{\partial \phi(y_m)}{\partial y_m} \end{pmatrix} \in M_m(K[t_1, \dots, t_m]),$$

where $M_m(K[t_1, \dots, t_m])$ is the associative algebra of $m \times m$ matrices with entries from $K[t_1, \dots, t_m]$. Let $\text{IE}(F_m)$ be the multiplicative semigroup of all endomorphisms of F_m which are identical modulo the commutator ideal F'_m . Let I_m be the identity $m \times m$ matrix and let S be the subspace of $M_m(K[t_1, \dots, t_m])$ defined by

$$S = \left\{ (f_{ij}) \in M_m(K[t_1, \dots, t_m]) \mid \sum_{i=1}^m t_i f_{ij} = 0, j = 1, \dots, m \right\}.$$

Clearly $I_m + S$ is a subsemigroup of the multiplicative group of $M_m(K[t_1, \dots, t_m])$. The completion with respect to the formal power series topology of $A_m \text{ wr } B_m$ is the semidirect sum $\widehat{C_m} \ltimes B_m$, where

$$\widehat{C_m} = \bigoplus_{i=1}^m a_i K[[t_1, \dots, t_m]].$$

If $\phi \in \text{IE}(\widehat{F_m})$, then $J(\phi) = I_m + (s_{ij})$, where $s_{ij} \in K[[t_1, \dots, t_m]]$. It is easy to check that if $\phi, \psi \in \text{IE}(\widehat{F_m})$ then $J(\phi\psi) = J(\phi)J(\psi)$. The following proposition is well known, see e.g. [BD].

Proposition 3. *The map $J : \text{IE}(F_m) \rightarrow I_m + S$ defined by $\phi \rightarrow J(\phi)$ is an isomorphism of the semigroups $\text{IE}(F_m)$ and $I_m + S$. It extends to an isomorphism between the group $\text{IE}(\widehat{F_m}) = \text{IA}(\widehat{F_m})$ of continuous IA-automorphisms of $\widehat{F_m}$ and the multiplicative group $I_m + \widehat{S}$.*

2. MAIN RESULTS

In this section we give a formula for the multiplication of inner automorphisms of $\widehat{F_m}$. We also find the explicit form of the Jacobian matrix of the inner automorphisms of $\widehat{F_m}$ and of the coset representatives of the continuous outer automorphisms in $\text{IOut}(\widehat{F_m})$. Finally we transfer the obtained results to the algebra $L_{m,c}$ and obtain the description of $\text{Inn}(L_{m,c})$ and $\text{IOut}(L_{m,c})$.

Let $\text{Inn}(\widehat{F_m})$ denote the set of all inner automorphisms of $\widehat{F_m}$ which are of the form $\exp(\text{adu})$, $u \in \widehat{F_m}$. Our first goal is to give a multiplication rule for $\text{Inn}(\widehat{F_m})$. For $u \in \widehat{F_m}$ we fix the notation $u = \overline{u} + u_0$, where $\overline{u}$ is the linear component of u and $u_0 \in F'_m$.

Theorem 4. *Let $u, v \in \widehat{F_m}$. Then the solution $w = w(u, v) \in \widehat{F_m}$ of the equation $\exp(\text{adu}) \cdot \exp(\text{adv}) = \exp(\text{ad}w)$ is*

$$w = H(\overline{u}, \overline{v}) + u_0(1 + (\text{ad}\overline{v})c(\text{ad}\overline{u}, \text{ad}\overline{v})) + v_0(1 - (\text{ad}\overline{u})c(\text{ad}\overline{u}, \text{ad}\overline{v})),$$

where

$$H(\overline{u}, \overline{v}) = \overline{u} + \overline{v} + [\overline{u}, \overline{v}]c(\text{ad}\overline{u}, \text{ad}\overline{v}),$$

$$c(t_1, t_2) = \frac{e^{t_2} h(t_1) - h(t_2)}{e^{t_1+t_2} - 1}, \quad h(t) = \frac{e^t - 1}{t}.$$

Proof. The adjoint operator $\text{ad} : \widehat{F_m} \rightarrow \text{End}_K \widehat{F_m}$ is a Lie algebra homomorphism. The algebra $\text{ad}(\widehat{F_m})$ is metabelian, as a homomorphic image of $\widehat{F_m}$. Hence we may apply the formula of Gerritzen from Proposition 1 for $x = \text{adu}$, $y = \text{adv}$, $z = \text{ad}w$. Therefore

$$\begin{aligned} \text{ad}w &= \text{adu} + \text{adv} + [\text{adu}, \text{adv}]c(\text{ad}(\text{adu}), \text{ad}(\text{adv})) \\ &= \text{adu} + \text{adv} + \text{ad}([u, v]c(\text{adu}, \text{adv})). \end{aligned}$$

Since the algebra $\widehat{F_m}$ has no centre, the adjoint representation is faithful and

$$w = u + v + [u, v]c(\text{adu}, \text{adv}).$$

The metabelian law gives that

$$\begin{aligned} [u, v]c(\text{adu}, \text{adv}) &= [\bar{u} + u_0, \bar{v} + v_0]c(\text{ad}(\bar{u} + u_0), \text{ad}(\bar{v} + v_0)) \\ &= ([\bar{u}, \bar{v}] + [u_0, \bar{v}] - [v_0, \bar{u}])c(\text{ad}\bar{u}, \text{ad}\bar{v}), \\ w &= (\bar{u} + u_0) + (\bar{v} + v_0) + ([\bar{u}, \bar{v}] + [u_0, \bar{v}] - [v_0, \bar{u}])c(\text{ad}\bar{u}, \text{ad}\bar{v}) \\ &= H(\bar{u}, \bar{v}) + u_0(1 + (\text{ad}\bar{v})c(\text{ad}\bar{u}, \text{ad}\bar{v})) + v_0(1 - (\text{ad}\bar{u})c(\text{ad}\bar{u}, \text{ad}\bar{v})). \end{aligned}$$

□

Our next objective is to give the explicit form of the Jacobian matrix of the inner automorphisms of $\widehat{F_m}$.

Theorem 5. Let $u = \bar{u} + u_0 \in \widehat{F_m}$, where

$$\bar{u} = \sum_{r=1}^m c_r y_r, \quad c_r \in K, r = 1, \dots, m,$$

is the linear component of u and

$$u_0 = \sum_{p>q} [y_p, y_q] h_{pq}(\text{ad}y_q, \dots, \text{ad}y_m).$$

Then

$$J(\exp(\text{adu})) = J(\exp(\text{ad}\bar{u})) + D_0 T = I_m + (\bar{D} + D_0)T,$$

$$\bar{D} = \left(\frac{\partial [y_j, \bar{u}]}{\partial y_i} \right), \quad D_0 = \left(\frac{\partial [y_j, u_0]}{\partial y_i} \right),$$

$$T = \sum_{k \geq 0} \frac{t^k}{(k+1)!}, \quad t = \sum_{r=1}^m c_r t_r.$$

More precisely

$$\begin{aligned} \bar{D} &= \begin{pmatrix} c_2 t_2 + c_3 t_3 + \dots + c_m t_m & -c_1 t_2 & \dots & -c_1 t_m \\ -c_2 t_1 & c_1 t_1 + c_3 t_3 + \dots + c_m t_m & \dots & -c_2 t_m \\ \vdots & \vdots & \ddots & \vdots \\ -c_m t_1 & -c_m t_2 & \dots & \sum_{k=1}^{m-1} c_k t_k \end{pmatrix}, \\ D_0 &= \begin{pmatrix} -t_1 f_1 & -t_2 f_1 & \dots & -t_m f_1 \\ -t_1 f_2 & -t_2 f_2 & \dots & -t_m f_2 \\ \vdots & \vdots & \ddots & \vdots \\ -t_1 f_m & -t_2 f_m & \dots & -t_m f_m \end{pmatrix}, \end{aligned}$$

where

$$\begin{aligned} f_i &= \sum_{p>q} \frac{\partial ([y_p, y_q] h_{pq}(\text{ad} y_q, \dots, \text{ad} y_m))}{\partial y_i} \\ &= \sum_{q=1}^{i-1} t_q h_{iq}(t_q, \dots, t_m) - \sum_{p=i+1}^m t_p h_{pi}(t_i, \dots, t_m). \end{aligned}$$

Proof. If $u = \bar{u} + u_0$, then

$$\exp(\text{adu})(y_j) = y_j + [y_j, u] \sum_{k \geq 0} \frac{(\text{ad} \bar{u})^k}{(k+1)!}, \quad j = 1, \dots, m,$$

because $u_0 \in \widehat{F}_m$ and adu_0 acts trivially on the commutator ideal of $\widehat{F}_m$. Hence

$$\begin{aligned} \exp(\text{adu})(y_j) &= y_j + [y_j, \bar{u} + u_0] \sum_{k \geq 0} \frac{(\text{ad} \bar{u})^k}{(k+1)!} \\ &= y_j + [y_j, \bar{u}] \sum_{k \geq 0} \frac{(\text{ad} \bar{u})^k}{(k+1)!} + [y_j, u_0] \sum_{k \geq 0} \frac{(\text{ad} \bar{u})^k}{(k+1)!} \\ &= \exp(\text{ad} \bar{u})(y_j) + [y_j, u_0] \sum_{k \geq 0} \frac{(\text{ad} \bar{u})^k}{(k+1)!}. \end{aligned}$$

Easy calculations give

$$\begin{aligned} \frac{\partial \exp(\text{ad} \bar{u})(y_j)}{\partial y_i} &= \delta_{ij} + \frac{\partial [y_j, \bar{u}]}{\partial y_i} \sum_{k \geq 0} \frac{t^k}{(k+1)!}, \\ \frac{\partial}{\partial y_i} [y_j, u_0] \sum_{k \geq 0} \frac{(\text{ad} \bar{u})^k}{(k+1)!} &= \frac{\partial [y_j, u_0]}{\partial y_i} \sum_{k \geq 0} \frac{t^k}{(k+1)!}, \\ i, j &= 1, \dots, m, \quad t = \sum_{p=1}^m c_p t_p, \end{aligned}$$

where δ_{ij} is Kronecker symbol. This gives the expression

$$J(\exp(\text{adu})) = I_m + (\bar{D} + D_0)T, \quad T = \sum_{k \geq 0} \frac{t^k}{(k+1)!},$$

$$\bar{D} = \left(\frac{\partial [y_j, \bar{u}]}{\partial y_i} \right), \quad D_0 = \left(\frac{\partial [y_j, u_0]}{\partial y_i} \right).$$

Since

$$\frac{\partial [y_j, \bar{u}]}{\partial y_i} = \frac{\partial}{\partial y_i} \left[y_j, \sum_{r=1}^m c_r y_r \right] = \frac{\partial}{\partial y_i} \sum_{r \neq j} c_r [y_j, y_r] = \begin{cases} \sum_{r \neq j} c_r t_r & i = j, \\ -c_i t_j & i \neq j, \end{cases}$$

we obtain the desired form of the matrix $\bar{D}$. Further,

$$[y_j, u_0] = - \left(\sum_{p>q} [y_p, y_q] h_{pq}(\text{ad} y_q, \dots, \text{ad} y_m) \right) \text{ad} y_j$$

$$\frac{\partial [y_j, u_0]}{\partial y_i} = -t_j \sum_{p>q} \frac{\partial [y_p, y_q]}{\partial y_i} h_{pq}(t_q, \dots, t_m), \quad \frac{\partial [y_p, y_q]}{\partial y_i} = \begin{cases} t_q & p = i, \\ -t_p & q = i, \\ 0 & p, q \neq i, \end{cases}$$

$$\frac{\partial[y_j, u_0]}{\partial y_i} = -t_j f_i(t_1, \dots, t_m),$$

$$f_i(t_1, \dots, t_m) = \sum_{q=1}^{i-1} t_q h_{iq}(t_q, \dots, t_m) - \sum_{p=i+1}^m t_p h_{pi}(t_i, \dots, t_m).$$

In this way we obtain the explicit form of the matrix D_0 . $\square$

Now we shall find the coset representatives of the normal subgroup $\text{Inn}(\widehat{F_m})$ of the group $\text{IA}(\widehat{F_m})$ of IA-automorphisms $\widehat{F_m}$, i.e., we shall find a set of IA-automorphisms θ of $\widehat{F_m}$ such that the factor group $\text{IOut}(\widehat{F_m}) = \text{IA}(\widehat{F_m})/\text{Inn}(\widehat{F_m})$ of the outer IA-automorphisms of $\widehat{F_m}$ is presented as the disjoint union of the cosets $\text{Inn}(\widehat{F_m})\theta$.

Theorem 6. *Let Θ be the set of automorphisms θ of $\widehat{F_m}$ with Jacobian matrix of the form*

$$J(\theta) = I_m + \begin{pmatrix} s(t_2, \dots, t_m) & f_{12} & \cdots & f_{1m} \\ t_1 q_2(t_2, t_3, \dots, t_m) + r_2(t_2, \dots, t_m) & f_{22} & \cdots & f_{2m} \\ t_1 q_3(t_3, \dots, t_m) + r_3(t_2, \dots, t_m) & f_{32} & \cdots & f_{3m} \\ \vdots & \vdots & \ddots & \vdots \\ t_1 q_m(t_m) + r_m(t_2, \dots, t_m) & f_{m2} & \cdots & f_{mm} \end{pmatrix},$$

where $s, q_i, r_i, f_{ij} \in K[[t_1, \dots, t_m]]$ are formal power series without constant terms and satisfy the conditions

$$s + \sum_{i=2}^m t_i q_i = 0, \quad \sum_{i=2}^m t_i r_i = 0, \quad \sum_{i=1}^m t_i f_{ij} = 0, \quad j = 2, \dots, m,$$

$r_i = r_i(t_2, \dots, t_m)$, $i = 2, \dots, m$, does not depend on t_1 , $q_i(t_i, \dots, t_m)$, $i = 2, \dots, m$, does not depend on $t_1, \dots, t_{i-1}$ and f_{12} does not contain a summand dt_2 , $d \in K$. Then Θ consists of coset representatives of the subgroup $\text{Inn}(\widehat{F_m})$ of the group $\text{IA}(\widehat{F_m})$ and $\text{IOut}(\widehat{F_m})$ is a disjoint union of the cosets $\text{Inn}(\widehat{F_m})\theta$, $\theta \in \Theta$.

Proof. Let $A = I_m + (f_{ij}) \in I_m + \widehat{S}$,

$$f_{11} = s, \quad f_{i1} = t_1 q_i + r_i, \quad i = 2, \dots, m,$$

be an $m \times m$ matrix satisfying the conditions of the theorem. The equation

$$s + \sum_{i=2}^m t_i q_i = 0$$

implies that

$$t_1 s + \sum_{i=2}^m t_i (t_1 q_i) = 0.$$

Hence By Lemma 2 gives that there exists an f_1 in the commutator ideal of $\widehat{F_m}$ such that

$$\frac{\partial f_1}{\partial y_1} = s, \quad \frac{\partial f_1}{\partial y_i} = t_1 q_i, \quad i = 2, \dots, m.$$

Similarly, the conditions

$$\sum_{i=2}^m t_i r_i = 0, \quad \sum_{i=1}^m t_i f_{ij} = 0, \quad j = 2, \dots, m,$$

imply that there exist $f'_1, f_j, j = 2, \dots, m$, in $\widehat{F'_m}$ with

$$\begin{aligned} \frac{\partial f'_1}{\partial y_1} &= 0, \quad \frac{\partial f'_1}{\partial y_i} = r_i, \quad i = 2, \dots, m, \\ \frac{\partial f_j}{\partial y_i} &= f_{ij}, \quad i = 1, \dots, j, \quad j = 2, \dots, m. \end{aligned}$$

This means that A is the Jacobian matrix of a certain IA-automorphism of $\widehat{F'_m}$.

Now we shall show that for any $\psi \in \text{IA}(\widehat{F'_m})$ there exists an inner automorphism $\phi = \exp(\text{adu}) \in \text{Inn}(\widehat{F'_m})$ and an automorphism θ in Θ such that $\psi = \exp(\text{adu}) \cdot \theta$. Let ψ be an arbitrary element of $\text{IA}(\widehat{F'_m})$ and let

$$\psi(y_1) = y_1 + \sum_{k>l} [y_k, y_l] f_{kl}(\text{ady}_l, \dots, \text{ady}_m),$$

$$\psi(y_2) = y_2 + \sum_{k>l} [y_k, y_l] g_{kl}(\text{ady}_l, \dots, \text{ady}_m),$$

where $f_{kl} = f_{kl}(t_1, \dots, t_m), g_{kl} = g_{kl}(t_1, \dots, t_m) \in K[[t_1, \dots, t_m]]$. Let us denote the $m \times 2$ matrix consisting of the first two columns of $J(\psi)$ by $J(\psi)_2$. Then $J(\psi)_2$ is of the form

$$J(\psi)_2 = \begin{pmatrix} 1 - t_2 f_{21} - t_3 f_{31} - \dots - t_m f_{m1} & -t_2 g_{21} - t_3 g_{31} - \dots - t_m g_{m1} \\ t_1 f_{21} - t_3 f_{32} - \dots - t_m f_{m2} & 1 + t_1 g_{21} - t_3 g_{32} - \dots - t_m g_{m2} \\ t_1 f_{31} + t_2 f_{32} - \dots - t_m f_{m3} & * \\ \vdots & \vdots \\ t_1 f_{m1} + \dots + t_{(m-1)} f_{m(m-1)} & * \end{pmatrix},$$

where we have denoted by $*$ the corresponding entries of the second column of the Jacobian matrix of ψ . Let

$$c_1 = -g_{21}(0, \dots, 0), \quad c_k = f_{k1}(0, \dots, 0), \quad k = 2, \dots, m,$$

and let us define

$$\phi_0 = \exp(\text{ad}\bar{u}), \quad \bar{u} = \sum_{i=1}^m c_i y_i.$$

Then

$$J(\phi_0) = I_m + \begin{pmatrix} \sum_{i \neq 1} c_i t_i & -c_1 t_2 & \dots & -c_1 t_m \\ -c_2 t_1 & \sum_{i \neq 2} c_i t_i & \dots & -c_2 t_m \\ -c_3 t_1 & -c_3 t_2 & \dots & -c_3 t_m \\ \vdots & \vdots & \ddots & \vdots \\ -c_m t_1 & -c_m t_2 & \dots & \sum_{i \neq m} c_i t_i \end{pmatrix} + B_0,$$

where the entries of the $m \times m$ matrix B_0 are elements of $K[[t_1, \dots, t_m]]$ which do not contain constant and linear terms. Hence

$$J(\phi_0 \psi)_2 = \begin{pmatrix} 1 + t_1 s_1(t_1, \dots, t_m) + s_2(t_2, \dots, t_m) & g(\widehat{t_2}) \\ t_1^2 p_2(t_1, \dots, t_m) + t_1 q_2(t_2, \dots, t_m) + r_2(t_2, \dots, t_m) & * \\ t_1^2 p_3(t_1, \dots, t_m) + t_1 q_3(t_2, \dots, t_m) + r_3(t_2, \dots, t_m) & * \\ \vdots & \vdots \\ t_1^2 p_m(t_1, \dots, t_m) + t_1 q_m(t_2, \dots, t_m) + r_m(t_2, \dots, t_m) & * \end{pmatrix},$$

where $t_1 s_1(t_1, \dots, t_m)$ and $s_2(t_2, \dots, t_m)$ have no linear terms, and $g(\widehat{t_2})$ does not contain a summand of the form dt_2 , $d \in K$. In the first column of $J(\phi_0\psi)_2$ we have collected the components $t_1^2 p_i$ divisible by t_1^2 , then the components $t_1 q_i$ divisible by t_1 only (but not by t_1^2) and finally the components r_i which do not depend on t_1 , $i = 2, \dots, m$. By Lemma 2 we obtain

$$t_1^2(s_1 + t_2 p_2 + \dots + t_m p_m) = 0,$$

$$t_1(s_2 + t_2 q_2 + \dots + t_m q_m) = 0,$$

$$t_2 r_2 + \dots + t_m r_m = 0.$$

Let us define $T_s = \{t_s, \dots, t_m\}$ and rewrite $J(\phi_0\psi)_2$ as

$$J(\phi_0\psi)_2 = \begin{pmatrix} 1 - t_1 t_2 p_2 - \dots - t_1 t_m p_m - t_2 q_2 - \dots - t_m q_m & g(\widehat{t_2}) \\ t_1^2 p_2 + t_1 q_2(T_2) + r_2(T_2) & * \\ t_1^2 p_3 + t_1 q_3(T_2) + r_3(T_2) & * \\ \vdots & \vdots \\ t_1^2 p_m + t_1 q_m(T_2) + r_m(T_2) & * \end{pmatrix},$$

Now we define

$$\phi_1 = \exp(\text{adu}_1), \quad u_1 = \sum_{i=2}^m [y_i, y_1] p_i(\text{ady}_1, \dots, \text{ady}_m).$$

The Jacobian matrix of ϕ_1 has the form

$$J(\phi_1) = I_m + \begin{pmatrix} t_1 \sum_{i \neq 1} t_i p_i & t_2 \sum_{i \neq 1} t_i p_i & \dots & t_m \sum_{i \neq 1} t_i p_i \\ -t_1^2 p_2 & -t_1 t_2 p_2 & \dots & -t_1 t_m p_2 \\ \vdots & \vdots & \ddots & \vdots \\ -t_1^2 p_m & -t_1 t_2 p_m & \dots & -t_1 t_m p_m \end{pmatrix}.$$

The element u_1 belongs to the commutator ideal of $\widehat{F_m}$ and the linear operator adu_1 acts trivially on $\widehat{F'_m}$. Hence $\exp(\text{adu}_1)$ is the identity map restricted on $\widehat{F'_m}$. Since the automorphism $\phi_0\psi$ is IA, we obtain that

$$\phi_0\psi(y_j) \equiv y_j \pmod{\widehat{F'_m}}, \quad \phi_1(\phi_0\psi(y_j)) = \phi_0\psi(y_j) + y_j \text{adu}_1.$$

Easy calculations give that

$$J(\phi_1\phi_0\psi)_2 = \begin{pmatrix} 1 - t_2 p_2 - \dots - t_m p_m & g(\widehat{t_2}) \\ t_1 q_2(T_2) + r_2(T_2) & * \\ t_1 q_3(T_2) + r_3(T_2) & * \\ \vdots & \vdots \\ t_1 q_m(T_2) + r_m(T_2) & * \end{pmatrix}.$$

Now we write $q_i(T_2)$ in the form

$$q_i(T_2) = t_2 q'_i(T_2) + q''_i(T_3), \quad i = 3, \dots, m,$$

and define

$$\phi_2 = \exp(\text{adu}_2), \quad u_2 = \sum_{i=3}^m [y_i, y_2] q'_i(\text{ady}_2, \dots, \text{ady}_m).$$

Then we obtain that

$$J(\phi_2\phi_1\phi_0\psi)_2 = \begin{pmatrix} 1 - t_2p_2 - \cdots - t_mp_m & g(\widehat{t_2}) \\ t_1Q_2(T_2) + r_2(T_2) & * \\ t_1q_3''(T_3) + r_3(T_2) & * \\ \vdots & \vdots \\ t_1q_m''(T_3) + r_m(T_2) & * \end{pmatrix},$$

$$Q_2(T_2) = q_2(T_2) - \sum_{i=3}^m t_i q_i'(T_2).$$

Repeating this process we construct inner automorphisms $\phi_3, \dots, \phi_{m-1}$ such that

$$\begin{aligned} \theta &= \phi_{m-1} \cdots \phi_2 \phi_1 \phi_0 \psi, \\ J(\phi_{m-1} \cdots \phi_2 \phi_1 \phi_0 \psi)_2 &= \begin{pmatrix} 1 + s(T_2) & g(\widehat{t_2}) \\ t_1Q_2(T_2) + r_2(T_2) & * \\ t_1Q_3(T_3) + r_3(T_2) & * \\ \vdots & \vdots \\ t_1Q_m(T_m) + r_m(T_2) & * \end{pmatrix}, \\ s(T_2) &= -t_2p_2(T_2) - \cdots - t_mp_m(T_2). \end{aligned}$$

Hence, starting from an arbitrary coset of IA-automorphisms $\text{Inn}(\widehat{F_m})\psi$, we found that it contains an automorphism $\theta \in \Theta$ with Jacobian matrix prescribed in the theorem. Now, let θ_1 and θ_2 be two different automorphisms in Θ with $\text{Inn}(\widehat{F_m})\theta_1 = \text{Inn}(\widehat{F_m})\theta_2$. Hence, there exists a nonzero element $u \in \widehat{F_m}$ such that $\theta_1 = \exp(\text{adu})\theta_2$. Direct calculations show that this is in contradiction with the form of $J(\theta_1)$. $\square$

Example 7. When $m = 2$ the results of Theorems 5 and 6 have the following simple form. If $u = \overline{u} + u_0$,

$$\overline{u} = c_1y_1 + c_2y_2, \quad u_0 = [y_2, y_1]h(\text{ady}_1, \text{ady}_2), \quad h(t_1, t_2) \in K[[t_1, t_2]],$$

then the Jacobian matrix of the inner automorphism $\exp(\text{adu})$ is

$$J(\exp(\text{adu})) = I_2 + \begin{pmatrix} (c_2 + t_1h)t_2 & (-c_1 + t_2h)t_2 \\ -(c_2 + t_1h)t_1 & -(-c_1 + t_2h)t_1 \end{pmatrix} \sum_{k \geq 0} \frac{(c_1t_1 + c_2t_2)^k}{(k+1)!}.$$

The Jacobian matrix of the outer automorphism $\theta \in \Theta$ is

$$J(\theta) = \begin{pmatrix} t_2f_1(t_2) & t_2f_2(t_1, t_2) \\ -t_1f_1(t_2) & -t_1f_2(t_1, t_2) \end{pmatrix}, \quad f_2(0, 0) = 0.$$

Recall that the augmentation ideal ω of the polynomial algebra $K[t_1, \dots, t_m]$ consists of the polynomials without constant terms and its completion $\widehat{\omega}$ is the ideal of $K[[t_1, \dots, t_m]]$ of all formal power series without constant terms. The elements of the commutator ideal of the free metabelian nilpotent of class c Lie algebra $L_{m,c}$ are of the form

$$u_0 = \sum_{p > q} [y_p, y_q] h_{pq}(\text{ady}_q, \dots, \text{ady}_m),$$

where $h_{pq}(t_q, \dots, t_m)$ belongs to the factor algebra $K[t_1, \dots, t_m]/\omega^{c-1}$ and may be identified with a polynomial of degree $\leq c-2$. The partial derivative $\partial u / \partial x_i$ belongs to $K[t_1, \dots, t_m]/\omega^c$ and may be considered as a polynomial of degree $\leq c-1$.

Similarly, the Jacobian matrix of an endomorphism of $F_{m,c}$ is considered modulo the ideal $M_m(\omega^c)$ of $M_m(K[t_1, \dots, t_m])$.

As a consequence of our Theorems 5 and 6 for $\text{Inn}(\widehat{F}_m)$ and $\text{IOut}(\widehat{F}_m)$ we immediately obtain the description of the groups of inner and outer automorphisms of $L_{m,c}$. We shall give the results for the Jacobian matrices only. The multiplication rule for the group $\text{Inn}(L_{m,c})$ from Theorem 4 can be stated similarly.

Corollary 8. (i) Let $u = \bar{u} + u_0 \in L_{m,c}$, where

$$\bar{u} = \sum_{r=1}^m c_r y_r, \quad c_r \in K, r = 1, \dots, m,$$

is the linear component of u and

$$u_0 = \sum_{p>q} [y_p, y_q] h_{pq}(\text{ad} y_q, \dots, \text{ad} y_m), \quad h_{pq}(t_1, \dots, t_m) \in K[t_1, \dots, t_m]/\omega^{c-1}.$$

Then the Jacobian matrix of the inner automorphism $\exp(\text{adu})$ is

$$J(\exp(\text{adu})) = J(\exp(\text{ad}\bar{u})) + D_0 T \equiv I_m + (\bar{D} + D_0) T \pmod{M_m(\omega^c)},$$

$$\bar{D} = \left(\frac{\partial[y_j, \bar{u}]}{\partial y_i} \right), \quad D_0 \equiv \left(\frac{\partial[y_j, u_0]}{\partial y_i} \right) \pmod{M_m(\omega^c)},$$

$$T = \sum_{k=0}^{c-1} \frac{t^k}{(k+1)!}, \quad t = \sum_{r=1}^m c_r t_r.$$

(ii) The automorphisms with the following Jacobian matrices are coset representatives of the subgroup $\text{Inn}(L_{m,c})$ of the group $\text{IA}(L_{m,c})$:

$$J(\theta) = I_m + \begin{pmatrix} s(t_2, \dots, t_m) & f_{12} & \cdots & f_{1m} \\ t_1 q_2(t_2, t_3, \dots, t_m) + r_2(t_2, \dots, t_m) & f_{22} & \cdots & f_{2m} \\ t_1 q_3(t_3, \dots, t_m) + r_3(t_2, \dots, t_m) & f_{32} & \cdots & f_{3m} \\ \vdots & \vdots & \ddots & \vdots \\ t_1 q_m(t_m) + r_m(t_2, \dots, t_m) & f_{m2} & \cdots & f_{mm} \end{pmatrix},$$

where $s, q_i, r_i, f_{ij} \in \omega/\omega^c$, i.e., are polynomials of degree $\leq c-1$ without constant terms. They satisfy the conditions

$$s + \sum_{i=2}^m t_i q_i \equiv 0, \quad \sum_{i=2}^m t_i r_i \equiv 0, \quad \sum_{i=1}^m t_i f_{ij} \equiv 0 \pmod{\omega^{c+1}}, \quad j = 2, \dots, m,$$

$r_i = r_i(t_2, \dots, t_m)$, $i = 1, \dots, m$, does not depend on t_1 , $q_i(t_i, \dots, t_m)$, $i = 2, \dots, m$, does not depend on $t_1, \dots, t_{i-1}$ and f_{12} does not contain a summand dt_2 , $d \in K$.

ACKNOWLEDGEMENTS

The second named author is grateful to the Institute of Mathematics and Informatics of the Bulgarian Academy of Sciences for the creative atmosphere and the warm hospitality during his visit when most of this project was carried out.

REFERENCES

- [Ba] Yu.A. Bahturin, Identical Relations in Lie Algebras (Russian), “Nauka”, Moscow, 1985. Translation: VNU Science Press, Utrecht, 1987.
- [Bo] N. Bourbaki, Éléments de mathématique. Fasc. XXXVII: Groupes et algèbres de Lie. Chap. II: Algèbres de Lie libres. Chap. III: Groupes de Lie, Actualites scientifiques et industrielles 1349. Paris: Hermann, 1972. Translation: Elements of mathematics. Lie Groups and Lie Algebras. Chapters 1-3, Berlin etc.: Springer-Verlag, 1989.
- [BD] R.M. Bryant, V. Drensky, Dense subgroups of the automorphism groups of free algebras, Canad.J.Math. 45 (1993), 1135-1154.
- [G] L. Gerritzen, Taylor expansion of noncommutative power series with an application to the Hausdorff series, J. Reine Angew. Math. 556 (2003), 113-125.
- [Se] J.-P. Serre, Lie Algebras and Lie Groups, W. A. Benjamin Inc., New York-Amsterdam, 1965.
- [Sh] A.L. Shmel’kin, Wreath products of Lie algebras and their application in the theory of groups (Russian), Trudy Moskov. Mat. Obshch. 29 (1973), 247-260. Translation: Trans. Moscow Math. Soc. 29 (1973), 239-252.

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, 1113
SOFIA, BULGARIA

E-mail address: `drensky@math.bas.bg`

DEPARTMENT OF MATHEMATICS, ÇUKUROVA UNIVERSITY, 01330 BALCALI, ADANA, TURKEY

E-mail address: `sfindik@cu.edu.tr`